
# FV<sup>3</sup>-powered Global-to-Regional Modeling at GFDL and Beyond



Lucas Harris, S-J Lin, Shannon Rees, Linjiong Zhou, Rusty Benson, Jan-Huey Chen, Kun Gao, Andy Hazelton, Bill Stern, and the GFDL FV<sup>3</sup> team

Modeling, Analysis, Predictions, and Projections Webinar 13 March 2017







Engineered to Make a Difference

#### The GFDL FV<sup>3</sup> Team














S-J Lin, Team Leader
Rusty Benson, Lead Engineer
Morris Bender Princeton Univ.
Jan-Huey Chen UCAR
Xi Chen Princeton Univ.
Kun Gao Princeton Univ.

Lucas Harris NOAA/GFDL
Andy Hazelton Princeton Univ.
Zhi Liang NOAA/GFDL
Tim Marchok NOAA/GFDL
Matt Morin Engility
Bill Putman NASA/GSFC

Shannon Rees Engility
Bill Stern UCAR
Bao Xiang UCAR
Weiye Yao Princeton Univ.
Ming Zhao NOAA/GFDL
Linjiong Zhou Princeton Univ.

#### WHAT IS FV<sup>3</sup>? FV<sup>3</sup> IS:




• Fully finite volume! Flux divergences + vertical Lagrangian + integrated PGF with excellent advection, nonhydrostatic option, and rigorous thermodynamics

#### But...

FV<sup>3</sup> is just a dynamical core not a model itself


## Needs physics, land model and ocean to be a complete model


- Proven effective at all scales. Maintains the large-scale circulation while accurately representing mesoscale and cloud-scale
- Popular! FV<sup>3</sup> the global core of choice for NOAA and NASA; predecessor FV standard in CESM and GEOS-CHEM community models, with FV<sup>3</sup> coming soon



#### TWO-WAY NESTING IN FV<sup>3</sup>

- Simultaneous coupled, consistent global and regional solution. No waiting for a regional prediction!
- Different grids permit different parameterizations and timesteps; doesn't need a "compromise" for highresolution region
- Flexible! Great possibilities for combining nesting and stretching.









#### **NESTED-GRID HIRAM SUBSEASONAL FORECASTS**

- HiRAM 4: FV<sup>3</sup> and AM2 physics + GFDL six-category microphysics and UW/GFDL double-plume convection
- Subseasonal forecasts:
   Chen & Lin (2012, 2013) seasonal frozen SST anomalies
   methodology
  - 5-member perturbed-physics 30-d forecast ensemble from 1st of each month








### HIRAM SUBSEASONAL FORECASTS 2000-2015 NORTH ATLANTIC MAJOR HURRICANES


25 km Uniform

8 km Nested










#### **3-KM NESTED fvGFS**


- fvGFS: FV<sup>3</sup> coupled to GFS physics
  - Prototype and proof-of-concept for next-generation GFS and unified global-to-regional forecast model
  - Simple GFS microphysics replaced with GFDL six-category scheme
- CONUS nest to 3 km for severe storm forecasts
  - Atlantic nest to 2km for hurricane forecasts




#### SEVERE CONVECTION IN fvGFS

2013 MOORE OUTBREAK FORECAST INITIALIZED OOZ 18 MAY (72 HOUR LEAD)
TUNED SAS-SHALLOW AND PBL, NO DEEP CONVECTION


#### fvGFS Max Reflectivity




#### fvGFS CONUS Max Reflectivity




#### Observed Max Reflectivity



#### fvGFS 850 mb Updraft Helicity



#### Observed Composite Reflectivity 20120629 1800Z



fvGFS Composite Reflectivity (-4 hr)




## SEVERE CONVECTION IN fvGFS

2012 DERECHO FORECAST INITIALIZED 00Z 27 JUNE (72 HOUR LEAD)

TUNED SAS-SHALLOW AND PBL,
NO DEEP CONVECTION


fvGFS w2500, lowest-level winds, T




#### **HURRICANE MATTHEW**


## INITIALIZED 5 OCT 2016 DEEP CONVECTION ON TUNED SAS-SHALLOW AND PBL

Courtesy Andy Hazelton and Morris Bender






#### **GLOBAL CONVECTION-RESOLVING FORECASTING**



2009: NASA GEOS-5 at 3.5 km global resolution



2017: GFDL fvGFS at 3-km global resolution

45 mins/day with 18.5K processors

Courtesy S-J Lin and Shannon Rees

#### PLANS FOR GLOBAL-TO-REGIONAL MODELING

- fvGFS: Working towards real-time 3-km system for HWT Spring Experiment
  - Improvements to land model, microphysics, and forecast diagnostics
  - Longer term: considering Thompson MP, YSU PBL, UW/GFDL Convection, etc.
- HiRAM & AM4: Bring grid refinement into next GFDL model
  - Convection-permitting (and resolving?) seasonal and longer simulations
- NCEP & AOML FV<sup>3</sup>-powered models:
  - 3-km NAM physics
  - Moving nested grids for next-generation HWRF
- NASA: 1.5-km GEOS forecasts coming soon
  - 3-km, year-long global nature run planned

## ADDITIONAL SLIDES

## PLANS FOR CONVECTION-RESOLVING fvGFS APPLICATIONS


- Thompson, Ferrier, or M-G Microphysics (collaboration with OU-CAPS and EMC)
- YSU, EDMF, or M-Y-type PBL
- Higher-resolution land model inputs
- Two fvGFS variants for the 2017 HWT Spring Experiment:
  - GFDL: fvGFS with GFDL MP
  - OU-CAPS: fvGFS with Thompson MP
- FV<sup>3</sup>-powered NAM and HWRF planned at EMC and AOML
- Convection-resolving and convection-permitting GFDL HiRAM/AM4 to continue

fvGFS is still improving! Stay tuned for further updates.

Special thanks to collaborators and others at CAPS, EMC, SPC, NSSL, AOML, NASA, and Harvard U.


#### TWO-WAY NESTING IN FV<sup>3</sup>


- Correct inflow/outflow BCs "baked-in" by FV<sup>3</sup> upwinding fluxes
- Concurrent nesting permits both domains to be run simultaneously
- Two-way updating of winds, w, and temperature consistent with finite-volume discretization
- Trivial mass conservation by not updating the mass field





#### 2015 HINDCAST SKILL IN NESTED FVGFS





CONUS Precipitation Fractions Skill Score
12 km neighborhood for nested,
40 km for uniform

fvGFS 13 km fvGFS 3 km (no deep) fvGFS 3 km (w/ deep) GFS 13 km

